arXiv:1606.05906v1 [math.CV] 19 Jun 2016

PARTIAL SUMS OF THE NORMALIZED DINI FUNCTIONS

HALİT ORHAN[1], İBRAHİM AKTAŞ[2]

ABSTRACT. Let $(w_{\alpha,v})_m(z) = z + \sum_{n=1}^{m} a_n z^{n+1}$ be the sequence of partial sums of normalized Dini functions $w_{\alpha,v}(z) = z + \sum_{n=1}^{\infty} a_n z^{n+1}$ where $a_n = \frac{(-1)^n (2n+\alpha)}{\alpha 4^n n! (v+1)_n}$. The aim of the present paper is to obtain lower bounds for $\mathcal{R}\left\{\frac{w_{\alpha,v}(z)}{(w_{\alpha,v})_m(z)}\right\}$, $\mathcal{R}\left\{\frac{(w_{\alpha,v})_m(z)}{w_{\alpha,v}(z)}\right\}$, $\mathcal{R}\left\{\frac{w'_{\alpha,v}(z)}{(w_{\alpha,v})'_m(z)}\right\}$ and $\mathcal{R}\left\{\frac{(w_{\alpha,v})'_m(z)}{w'_{\alpha,v}(z)}\right\}$. Also we give a few geometric description regarding image domains of some functions.

1. INTRODUCTION

Let $\mathcal{A}$ denote the family of functions f of the form

$$f(z) = z + \sum_{k=1}^{\infty} a_k z^{k+1} \tag{1.1}$$

which are analytic in the open unit disk $\mathcal{U} = \{z \in \mathcal{C} : |z| < 1\}$ and satisfy the usual normalization condition $f(0) = f'(0) - 1 = 0$. Let $\mathcal{S}$ denote the subclass of $\mathcal{A}$ which are univalent in $\mathcal{U}$. Also let $\mathcal{S}^*$and $\mathcal{C}$ denote the subclasses of $\mathcal{A}$ consisting of functions which are starlike and convex in $\mathcal{U}$,respectively. These classes can characterize as follows:

$$f \in \mathcal{S}^* \Leftrightarrow \Re\left(\frac{zf'(z)}{f(z)}\right) > 0, \tag{1.2}$$

$$f \in \mathcal{C} \Leftrightarrow \Re\left(1 + \frac{zf''(z)}{f'(z)}\right) > 0. \tag{1.3}$$

Some special functions like Bessel functions play an important role in applied mathematics and physics. We know that the Bessel functions of the first kind J_v is defined by(see [2] and [12])

$$J_v(z) = \sum_{n=0}^{\infty} \frac{(-1)^n}{n!\Gamma(v+n+1)} \left(\frac{z}{2}\right)^{2n+v} \quad (z \in \mathcal{C}) \tag{1.4}$$

where $\Gamma(z)$ stands for Euler gamma function and it is a particular solution of the second-order linear homogeneous differential equation(see, for details, [2] and [12]):

$$z^2 y''(z) + z y'(z) + \left[z^2 - v^2\right] y(z) = 0 \tag{1.5}$$

where $v \in \mathcal{C}$.

Date: 17.05.2016.

2000 *Mathematics Subject Classification.* 33C10, 30C45.

Key words and phrases. Dini functions; Bessel function; Analytic function; Univalent, Subordination.

Now, we consider the function $w_{\alpha,v}:\mathcal{U}\to\mathcal{C}$, defined by(see [4])

$$w_{\alpha,v}(z) = \frac{2^v}{\alpha}\Gamma(v+1)z^{1-\frac{v}{2}}((\alpha-v)J_v(\sqrt{z})+\sqrt{z}J_v^{'}(\sqrt{z})) = \sum_{n=0}^{\infty}\frac{(-1)^n(2n+\alpha)\Gamma(v+1)z^{n+1}}{\alpha.4^n n!\Gamma(v+n+1)}. \tag{1.6}$$

On the other hand we know that there is following relation in between Pochhammer Symbol (or Appell) and Euler's gamma function:

$$(\mu)_n = \frac{\Gamma(\mu+n)}{\Gamma(\mu)} = \mu(\mu+1)...(\mu+n-1). \tag{1.7}$$

If we use the relation (1.7) we can obtain the following series representation for the functions $w_{\alpha,v}(z)$ given by (1.6):

$$w_{\alpha,v}(z) = z+\sum_{n=1}^{\infty}a_n z^{n+1} \tag{1.8}$$

where $a_n=\frac{(-1)^n(2n+\alpha)}{\alpha.4^n n!(v+1)_n}$.

In this study, we will investigate the ratio of a function of the form (1.8) to its sequence of partial sums $(w_{\alpha,v}(z))_m = z+\sum_{n=1}^{m}a_n z^{n+1}$ when the coefficients of $w_{\alpha,v}$ satisfy some conditions. We will obtain lower bounds for $\mathcal{R}\left\{\frac{w_{\alpha,v}(z)}{(w_{\alpha,v})_m(z)}\right\}$, $\mathcal{R}\left\{\frac{(w_{\alpha,v})_m(z)}{w_{\alpha,v}(z)}\right\}$, $\mathcal{R}\left\{\frac{w_{\alpha,v}^{'}(z)}{(w_{\alpha,v})_m^{'}(z)}\right\}$ and $\mathcal{R}\left\{\frac{(w_{\alpha,v})_m^{'}(z)}{w_{\alpha,v}^{'}(z)}\right\}$.

For interesting developments on the partial sums of some special functions and the some classes of analytic functions, the readers can browse to the works of Orhan and Yagmur [7], Çağlar and Deniz [6], Brickman et al. [5], Silverman [10], Owa et al. [8], Sheil-Small [9], Silvia [11], Aktaş and Orhan [1].

Lemma 1.1. *If $\alpha>0$ and $v>-\frac{7}{8}$, then the function*

$$w_{\alpha,v}:\mathcal{U}\to\mathcal{C}$$

given by (1.6) satisfies the follwing inequalities:

$$|w_{\alpha,v}(z)|\leq 1+\frac{32v+2\alpha+32}{(8v+7)\alpha}, \tag{1.9}$$

$$\left|w_{\alpha,v}^{'}(z)\right|\leq 1+\frac{256(2+\alpha)v^2+8(64+29\alpha)v+210\alpha+512}{(8v+7)^3\alpha}. \tag{1.10}$$

Proof. By using the well-known triangle inequality:

$$|z_1+z_2|\leq|z_1|+|z_2| \tag{1.11}$$

and the inequalities

$$(v+1)^n\leq(v+1)_n \text{ and } 2^{n-1}\leq n! \tag{1.12}$$

for $n \in \mathbb{N} = \{1, 2, ...\}$, we get

$$\begin{aligned} |w_{\alpha,v}(z)| &= \left| z + \sum_{n=1}^{\infty} \frac{(-1)^n (2n+\alpha)}{\alpha.4^n n!(v+1)_n} z^{n+1} \right| \leq 1 + \sum_{n=1}^{\infty} \frac{2n+\alpha}{\alpha.4^n 2^{n-1}(v+1)^n} \\ &= 1 + \frac{2}{4\alpha(v+1)} \sum_{n=1}^{\infty} \frac{n}{(8(v+1))^{n-1}} + \frac{1}{4(v+1)} \sum_{n=1}^{\infty} \left(\frac{1}{8(v+1)} \right)^{n-1} \\ &= 1 + \frac{32v + 2\alpha + 32}{(8v+7)\alpha}. \end{aligned}$$

If we consider inequality (1.10), then we have

$$\begin{aligned} \left| w'_{\alpha,v}(z) \right| &= \left| 1 + \sum_{n=1}^{\infty} \frac{(-1)^n (2n+\alpha)(n+1)}{\alpha.4^n n!(v+1)_n} z^n \right| \leq 1 + \sum_{n=1}^{\infty} \frac{2n^2 + (2+\alpha)n + \alpha}{\alpha.4^n 2^{n-1}(v+1)^n} \\ &= 1 + \frac{1}{(v+1)} \left\{ \frac{1}{2\alpha} \sum_{n=1}^{\infty} \frac{n^2}{(8(v+1))^{n-1}} + \frac{(2+\alpha)}{4\alpha} \sum_{n=1}^{\infty} \frac{n}{(8(v+1))^{n-1}} + \frac{1}{4} \sum_{n=1}^{\infty} \left(\frac{1}{8(v+1)} \right)^{n-1} \right\} \\ &= 1 + \frac{256(2+\alpha)v^2 + 8(64+29\alpha)v + 210\alpha + 512}{(8v+7)^3 \alpha}. \end{aligned}$$

□

2. Main Results

Theorem 2.1. *Let $\alpha > 0$, $v > -\frac{7}{8}$, the function $w_{\alpha,v} : \mathcal{U} \to \mathcal{C}$ defined by (1.8) and its sequence of partial sums $(w_{\alpha,v}(z))_m = z + \sum_{n=1}^{m} a_n z^{n+1}$. If the inequality*

$$8(\alpha - 4)v + 5\alpha - 32 \geq 0$$

is true, then we have the followings:

$$\mathcal{R}\left\{ \frac{w_{\alpha,v}(z)}{(w_{\alpha,v})_m(z)} \right\} \geq \frac{8(\alpha-4)v + 5\alpha - 32}{(8v+7)\alpha}, \tag{2.1}$$

$$\mathcal{R}\left\{ \frac{(w_{\alpha,v})_m(z)}{w_{\alpha,v}(z)} \right\} \geq \frac{(8v+7)\alpha}{(8v+7)\alpha + 32v + 2\alpha + 32}. \tag{2.2}$$

Proof. From the inequality (1.9) in the Lemma 1.1 we write that

$$1 + \sum_{n=1}^{\infty} |a_n| \leq 1 + \frac{32v + 2\alpha + 32}{(8v+7)\alpha}$$

which is equivalent to

$$\frac{(8v+7)\alpha}{32v + 2\alpha + 32} \sum_{n=1}^{\infty} |a_n| \leq 1,$$

where $a_n = \frac{(-1)^n (2n+\alpha)}{\alpha.4^n n!(v+1)_n}$.

Now we may write

$$\frac{(8v+7)\alpha}{32v + 2\alpha + 32} \left\{ \frac{w_{\alpha,v}(z)}{(w_{\alpha,v})_m(z)} - \left(1 - \frac{32v + 2\alpha + 32}{(8v+7)\alpha} \right) \right\} \tag{2.3}$$

$$= \frac{1+\sum_{n=1}^{m} a_n z^n + \frac{(8v+7)\alpha}{32v+2\alpha+32} \sum_{n=m+1}^{\infty} a_n z^n}{1+\sum_{n=1}^{m} a_n z^n}$$

$$= \frac{1+A(z)}{1+B(z)}.$$

Set $(1+A(z))/(1+B(z)) = (1+p(z))/(1-p(z))$, so that $p(z) = (A(z)-B(z))/(2+A(z)+B(z))$. Then

$$p(z) = \frac{\frac{(8v+7)\alpha}{32v+2\alpha+32} \sum_{n=m+1}^{\infty} a_n z^n}{2+2\sum_{n=1}^{m} a_n z^n + \frac{(8v+7)\alpha}{32v+2\alpha+32} \sum_{n=m+1}^{\infty} a_n z^n}$$

and

$$|p(z)| \leq \frac{\frac{(8v+7)\alpha}{32v+2\alpha+32} \sum_{n=m+1}^{\infty} |a_n|}{2-2\sum_{n=1}^{m} |a_n| - \frac{(8v+7)\alpha}{32v+2\alpha+32} \sum_{n=m+1}^{\infty} |a_n|}.$$

Now $|p(z)| \leq 1$ if and only if

$$\sum_{n=1}^{m} |a_n| + \frac{(8v+7)\alpha}{32v+2\alpha+32} \sum_{n=m+1}^{\infty} |a_n| \leq 1. \tag{2.4}$$

It suffices to show that the left hand side of (2.4) is bounded above by $\frac{(8v+7)\alpha}{32v+2\alpha+32} \sum_{n=1}^{\infty} |a_n|$, which is equivalent to

$$\frac{8(\alpha-4)v+5\alpha-32}{32v+2\alpha+32} \sum_{n=1}^{m} |a_n| \geq 0.$$

To prove the result (2.2), we write

$$\left(\frac{(8v+7)\alpha}{32v+2\alpha+32}+1\right)\left\{\frac{(w_{\alpha,v})_m(z)}{w_{\alpha,v}(z)} - \frac{(8v+7)\alpha}{(8v+7)\alpha+32v+2\alpha+32}\right\} \tag{2.5}$$

$$= \frac{1+\sum_{n=1}^{m} a_n z^n - \frac{(8v+7)\alpha}{32v+2\alpha+32} \sum_{n=m+1}^{\infty} a_n z^n}{1+\sum_{n=1}^{\infty} a_n z^n}$$

$$= \frac{1+p(z)}{1-p(z)}.$$

Then from (2.5) we get

$$p(z)=\frac{-\frac{8(\alpha+4)v+9\alpha+32}{32v+2\alpha+32}\sum\limits_{n=m+1}^{\infty}a_nz^n}{2+2\sum\limits_{n=1}^{m}a_nz^n-\frac{8(\alpha-4)v+5\alpha-32}{32v+2\alpha+32}\sum\limits_{n=m+1}^{\infty}a_nz^n}$$

and

$$|p(z)|\leq\frac{\frac{8(\alpha+4)v+9\alpha+32}{32v+2\alpha+32}\sum\limits_{n=m+1}^{\infty}|a_n|}{2-2\sum\limits_{n=1}^{m}|a_n|-\frac{8(\alpha-4)v+5\alpha-32}{32v+2\alpha+32}\sum\limits_{n=m+1}^{\infty}|a_n|}.$$

Now $|p(z)|\leq 1$ if and only if

$$\sum_{n=1}^{m}|a_n|+\frac{(8v+7)\,\alpha}{32v+2\alpha+32}\sum_{n=m+1}^{\infty}|a_n|\leq 1. \tag{2.6}$$

Since the left hand side of (2.6) is bounded above by $\frac{(8v+7)\alpha}{32v+2\alpha+32}\sum\limits_{n=1}^{\infty}|a_n|$, the proof is complated. □

Theorem 2.2. *Let $\alpha>0$, $v>-\frac{7}{8}$, the function $w_{\alpha,v}:\mathcal{U}\to\mathcal{C}$ defined by (1.8) and its sequence of partial sums $(w_{\alpha,v}(z))_m=z+\sum\limits_{n=1}^{m}a_nz^{n+1}$. If the inequality*

$$512\alpha v^3+64\,(17\alpha-8)\,v^2+16\,(59\alpha-32)\,v+133\alpha-512\geq 0$$

is true, then we have the followings:

$$\mathcal{R}\left\{\frac{w'_{\alpha,v}(z)}{(w_{\alpha,v})'_m(z)}\right\}\geq\frac{512\alpha v^3+64\,(17\alpha-8)\,v^2+16\,(59\alpha-32)\,v+133\alpha-512}{256\,(2+\alpha)\,v^2+8\,(64+29\alpha)\,v+210\alpha+512}, \tag{2.7}$$

$$\mathcal{R}\left\{\frac{(w_{\alpha,v})'_m(z)}{w'_{\alpha,v}(z)}\right\}\geq\frac{(8v+7)^3\,\alpha}{512\alpha v^3+64\,(25\alpha+8)\,v^2+128\,(11\alpha+4)\,v+553\alpha+512}. \tag{2.8}$$

Proof. From the inequality (1.10) in the Lemma 1.1 we write that

$$1+\sum_{n=1}^{\infty}(n+1)\,|a_n|\leq 1+\frac{256\,(2+\alpha)\,v^2+8\,(64+29\alpha)\,v+210\alpha+512}{(8v+7)^3\,\alpha}$$

which is equivalent to

$$\frac{(8v+7)^3\,\alpha}{256\,(2+\alpha)\,v^2+8\,(64+29\alpha)\,v+210\alpha+512}\sum_{n=1}^{\infty}(n+1)\,|a_n|\leq 1$$

where $a_n=\frac{(-1)^n(2n+\alpha)}{\alpha.4^n n!(v+1)_n}$.

Now we write

$$\begin{aligned}
&\frac{(8v+7)^3\alpha}{256(2+\alpha)v^2+8(64+29\alpha)v+210\alpha+512}\times\\
&\left\{\frac{w'_{\alpha,v}(z)}{(w_{\alpha,v})'_m(z)}-(1-\frac{256(2+\alpha)v^2+8(64+29\alpha)v+210\alpha+512}{(8v+7)^3\alpha})\right\}\\
=&\ \frac{1+\sum_{n=1}^{m}(n+1)a_nz^n+\frac{(8v+7)^3\alpha}{256(2+\alpha)v^2+8(64+29\alpha)v+210\alpha+512}\sum_{n=m+1}^{\infty}(n+1)a_nz^n}{1+\sum_{n=1}^{m}(n+1)a_nz^n}\\
=&\ \frac{1+p(z)}{1-p(z)}
\end{aligned}$$

where

$$|p(z)|\leq\frac{\frac{(8v+7)^3\alpha}{256(2+\alpha)v^2+8(64+29\alpha)v+210\alpha+512}\sum_{n=m+1}^{\infty}(n+1)|a_n|}{2-2\sum_{n=1}^{m}(n+1)|a_n|-\frac{(8v+7)^3\alpha}{256(2+\alpha)v^2+8(64+29\alpha)v+210\alpha+512}\sum_{n=m+1}^{\infty}(n+1)|a_n|}\leq 1.$$

The last inequality is equivalent to

$$\sum_{n=1}^{m}(n+1)|a_n|+\frac{(8v+7)^3\alpha}{256(2+\alpha)v^2+8(64+29\alpha)v+210\alpha+512}\sum_{n=m+1}^{\infty}(n+1)|a_n|\leq 1. \tag{2.9}$$

It suffices to show that the left hand side of (2.9) is bounded above by
$\frac{(8v+7)^3\alpha}{256(2+\alpha)v^2+8(64+29\alpha)v+210\alpha+512}\sum_{n=1}^{\infty}(n+1)|a_n|$, which is equivalent to

$$\frac{512\alpha v^3+64(17\alpha-8)v^2+16(59\alpha-32)+133\alpha-512}{256(2+\alpha)v^2+8(64+29\alpha)v+210\alpha+512}\sum_{n=1}^{m}(n+1)|a_n|\geq 0.$$

To prove the result (2.8), we write

$$\begin{aligned}
\frac{1+p(z)}{1-p(z)}=&\left(\frac{(8v+7)^3\alpha}{256(2+\alpha)v^2+8(64+29\alpha)v+210\alpha+512}+1\right)\times\\
&\left\{\frac{(w_{\alpha,v})'_m(z)}{w'_{\alpha,v}(z)}-\frac{(8v+7)^3\alpha}{512\alpha v^3+64(25\alpha+8)v^2+128(11\alpha+4)v+553\alpha+512}\right\}\\
=&\ \frac{1+\sum_{n=1}^{m}(n+1)a_nz^n-\frac{(8v+7)^3\alpha}{256(2+\alpha)v^2+8(64+29\alpha)v+210\alpha+512}\sum_{n=m+1}^{\infty}(n+1)a_nz^n}{1+\sum_{n=1}^{\infty}(n+1)a_nz^n}
\end{aligned}$$

where

$$|p(z)| \leq \frac{\frac{512\alpha v^3+64(25\alpha+8)v^2+128(11\alpha+4)v+553\alpha+512}{256(2+\alpha)v^2+8(64+29\alpha)v+210\alpha+512}\sum_{n=m+1}^{\infty}(n+1)|a_n|}{2-2\sum_{n=1}^{m}(n+1)|a_n|-\frac{512\alpha v^3+64(17\alpha-8)v^2+16(59\alpha-32)+133\alpha-512}{256(2+\alpha)v^2+8(64+29\alpha)v+210\alpha+512}\sum_{n=m+1}^{\infty}(n+1)|a_n|} \leq 1.$$

The last inequality is equivalent to

$$\sum_{n=1}^{m}(n+1)|a_n|+\frac{(8v+7)^3\alpha}{256(2+\alpha)v^2+8(64+29\alpha)v+210\alpha+512}\sum_{n=m+1}^{\infty}(n+1)|a_n| \leq 1. \tag{2.10}$$

Since the left hand side of (2.10) is bounded above by $\frac{(8v+7)^3\alpha}{256(2+\alpha)v^2+8(64+29\alpha)v+210\alpha+512}\sum_{n=1}^{\infty}(n+1)|a_n|$, the proof is complated. □

For some special cases of v it is known that(see [3])

$$J_{\frac{1}{2}}(z)=\sqrt{\frac{2}{\pi z}}\sin z \tag{2.11}$$

and

$$J_{\frac{3}{2}}(z)=\sqrt{\frac{2}{\pi z}}\left(\frac{\sin z}{z}-\cos z\right). \tag{2.12}$$

Now by using some special cases of α and v in Theorem 2.1 and Theorem 2.2, we can obtain following corollaries.

Corollary 2.1. *If we take $\alpha=1$, $v=\frac{1}{2}$ and $m=0$, we have $w_{1,\frac{1}{2}}(z)=\frac{\sqrt{z}}{2}\left\{\sin\sqrt{z}+\cos\sqrt{z}-\frac{\sin\sqrt{z}}{4}\right\}$, $w'_{1,\frac{1}{2}}(z)=\frac{1}{4\sqrt{z}}\left\{(1-\sqrt{z})\sin\sqrt{z}+\frac{(1+2\sqrt{z})\cos\sqrt{z}}{2}\right\}$, $\left(w_{1,\frac{1}{2}}\right)_0(z)=z$ and $\left(w_{1,\frac{1}{2}}\right)'_0(z)=1$. In view of Theorem 2.2 we get the following inequalities:*

$$\mathcal{R}\left\{\frac{1}{4\sqrt{z}}\left(\left(1-\sqrt{z}\right)\sin\sqrt{z}+\frac{(1+2\sqrt{z})\cos\sqrt{z}}{2}\right)\right\}\geq\frac{45}{1286},$$

$$\mathcal{R}\left\{\frac{8\sqrt{z}}{(1+2\sqrt{z})\cos\sqrt{z}-2\left(\sqrt{z}-1\right)\sin\sqrt{z}}\right\}\geq\frac{1331}{2617}.$$

Corollary 2.2. *If we take $\alpha=\frac{3}{2}$, $v=\frac{1}{2}$ and $m=0$, we have $w_{\frac{3}{2},\frac{1}{2}}(z)=\frac{\sqrt{z}}{3}(2\sin\sqrt{z}+\cos\sqrt{z})-\frac{\sin\sqrt{z}}{6}$, $w'_{\frac{3}{2},\frac{1}{2}}(z)=\frac{4\sin\sqrt{z}+\cos\sqrt{z}}{12\sqrt{z}}+\frac{2\cos\sqrt{z}-\sin\sqrt{z}}{6}$, $\left(w_{\frac{3}{2},\frac{1}{2}}\right)_0(z)=z$ and $\left(w_{\frac{3}{2},\frac{1}{2}}\right)'_0(z)=1$. In view of Theorem 2.2 we get the following inequalities:*

$$\mathcal{R}\left\{\frac{4\sin\sqrt{z}+\cos\sqrt{z}}{12\sqrt{z}}+\frac{2\cos\sqrt{z}-\sin\sqrt{z}}{6}\right\}\geq\frac{1031}{2962},$$

$$\mathcal{R}\left\{\frac{12\sqrt{z}}{(4\sqrt{z}+1)\cos\sqrt{z}-2\left(\sqrt{z}-2\right)\sin\sqrt{z}}\right\}\geq\frac{3993}{6955}.$$

Corollary 2.3. *If we take* $\alpha = 5$, $v = \frac{3}{2}$ *and* $m = 0$, *we have* $w_{5,\frac{3}{2}}(z) = \frac{21}{10}\left(\frac{\sin\sqrt{z}}{\sqrt{z}} - \cos\sqrt{z}\right) + \frac{9}{20}\left(\frac{\cos\sqrt{z}}{\sqrt{z}} - \frac{\sin\sqrt{z}}{z}\right) + \frac{3}{10}\sin\sqrt{z}$, $w'_{5,\frac{3}{2}}(z) = \frac{21}{20}\left(\frac{\cos\sqrt{z}}{z} - \frac{\sin\sqrt{z}}{z\sqrt{z}} + \frac{\sin\sqrt{z}}{\sqrt{z}}\right) + \frac{9}{20}\left(\frac{\sin\sqrt{z}}{z^2} - \frac{\cos\sqrt{z}}{z\sqrt{z}}\right) - \frac{9}{40}\frac{\sin\sqrt{z}}{z} + \frac{3}{20}\frac{\cos\sqrt{z}}{\sqrt{z}}$, $\left(w_{5,\frac{3}{2}}\right)_0(z) = z$ *and* $\left(w_{5,\frac{3}{2}}\right)'_0(z) = 1$. *In view of Theorem 2.1 we get the following inequalities:*

$$\mathcal{R}\left\{\frac{14z\left(\sin\sqrt{z} - \sqrt{z}\cos\sqrt{z}\right) + (2z-3)\sqrt{z}\sin\sqrt{z} + 3z\cos\sqrt{z}}{z^{5/2}}\right\} \geq \frac{20}{57},$$

$$\mathcal{R}\left\{\frac{z^{5/2}}{14z\left(\sin\sqrt{z} - \sqrt{z}\cos\sqrt{z}\right) + (2z-3)\sqrt{z}\sin\sqrt{z} + 3z\cos\sqrt{z}}\right\} \geq \frac{57}{740}.$$

3. Illustrative examples and image domains

In this section, illustrative examples along with the geometrical descriptions of the image domains of the unit disk by the ratio of normalized Dini function to its sequence of partial sums or the the ratio of its sequence of partial sums to the function which we considered in our corollaries in section 2, have been given in following figures.

Example 3.1. The image domains of $f_1(z) = \frac{1}{4\sqrt{z}}\left((1-\sqrt{z})\sin\sqrt{z} + \frac{(1+2\sqrt{z})\cos\sqrt{z}}{2}\right)$ and $f_2(z) = \frac{8\sqrt{z}}{(1+2\sqrt{z})\cos\sqrt{z} - 2(\sqrt{z}-1)\sin\sqrt{z}}$ are shown in Figure 1, while the image domains of $f_3(z) = \frac{4\sin\sqrt{z}+\cos\sqrt{z}}{12\sqrt{z}} + \frac{2\cos\sqrt{z}-\sin\sqrt{z}}{6}$ and $f_4(z) = \frac{12\sqrt{z}}{(4\sqrt{z}+1)\cos\sqrt{z} - 2(\sqrt{z}-2)\sin\sqrt{z}}$ are shown in Figure 2.

Figure 1 *Figure* 2

Example 3.2. We have the image domains of $f_5(z) = \frac{14z\left(\sin\sqrt{z}-\sqrt{z}\cos\sqrt{z}\right)+(2z-3)\sqrt{z}\sin\sqrt{z}+3z\cos\sqrt{z}}{z^{5/2}}$ and $f_6(z) = \frac{z^{5/2}}{14z\left(\sin\sqrt{z}-\sqrt{z}\cos\sqrt{z}\right)+(2z-3)\sqrt{z}\sin\sqrt{z}+3z\cos\sqrt{z}}$ in Figure 3 and Figure 4, respectively.

Figure 3 *Figure* 4

References

[1] Aktaş I, Orhan H. Distortion bounds for a new subclass of analytic functions and their partial sums. Bulletin of the Transilvania University of Braşov Series III 2015, 57 (2): 1-12.

[2] Baricz A. Geometric properties of generilazed Bessel function. Publ. Math. Debrecen 2008, 73, 1-2: 155-178.

[3] Baricz A. Generalized Bessel Functions of the First Kind. Lecture Notes in Mathematics Vol. 1994, Berlin, Springer-Verlag, 2010.

[4] Baricz A, Deniz E, Yağmur N. Close-to-convexity of normalized Dini functions. Math. Nachr.2016: 1-6.

[5] Brickman L, Hallenbeck DJ, Macgragor TH, Wilken D. Convex hulls and extreme points of families of starlike and convex mappings. Trans. Amer. Math. Soc. 1973, 185: 413–428.

[6] Çağlar M, Deniz E. Partial sums of the normalized Lommel functions. Math. Ineq. & Appl. 2015, 18 (3): 1189-1199.

[7] Orhan H, Yağmur N. Partial sums of Generilazed Bessel Functions. Journal of Mathematical Inequalities 2014,8 (4): 863-877.

[8] Owa S, Srivastava HM, Saito N. Partial sums of certain classes of analytic functions. Int. J. Comput. Math. 2004, 81, (10): 1239–1256.

[9] Sheil-Small T. A note on partial sums of convex schlicht functions. Bull. London Math. Soc. 1970, 2: 165–168.

[10] Silverman H. Partial sums of starlike and convex functions. J. Math. Anal. Appl. 1997, 209: 221–227.

[11] Silvia EM. On partial sums of convex functions of order α. Houston J. Math. 1985, 11: 397–404.

[12] Watson GN. A Tretaise on the Theory of Bessel Functions. Second edition, London and New York, Cambridge University Press, 1944.

[1]Department of Mathematics, Faculty of Science, Atatürk University, Erzurum, 25240, Turkey.
E-mail address: orhanhalit607@gmail.com

[2]Corresponding author, Department of Mathematical Engineering, Faculty of Engineering and Natural Science, Gümüşhane University, Gümüşhane, 29100, Turkey.
E-mail address: aktasibrahim38@gmail.com

This figure "f1-f2.png" is available in "png" format from:

http://arxiv.org/ps/1606.05906v1

This figure "f3-f4.png" is available in "png" format from:

http://arxiv.org/ps/1606.05906v1

This figure "f5.png" is available in "png" format from:

http://arxiv.org/ps/1606.05906v1

This figure "f6.png" is available in "png" format from:

http://arxiv.org/ps/1606.05906v1